

## Algebraic

- $a^2 - b^2 = (a - b)(a + b)$
- $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$
- Quadratic Formula:  $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

## Geometric

- Area of Circle:  $\pi r^2$
- Circumference of Circle:  $2\pi r$
- Circle with center  $(h, k)$  and radius  $r$ :  

$$(x - h)^2 + (y - k)^2 = r^2$$
- Distance from  $(x_1, y_1)$  to  $(x_2, y_2)$ :  

$$\sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$
- Area of Triangle:  $\frac{1}{2}bh$
- $\sin \theta = \frac{\text{opposite leg}}{\text{hypotenuse}}$
- $\cos \theta = \frac{\text{adjacent leg}}{\text{hypotenuse}}$
- $\tan \theta = \frac{\text{opposite leg}}{\text{adjacent leg}}$
- If  $\triangle ABC$  is similar to  $\triangle DEF$  then

$$\frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$$

- Volume of Sphere:  $\frac{4}{3}\pi r^3$
- Surface Area of Sphere:  $4\pi r^2$
- Volume of Cylinder/Prism: (height)(area of base)
- Volume of Cone/Pyramid:  $\frac{1}{3}(\text{height})(\text{area of base})$

## Theorems

- (IVT) If  $f$  is continuous on  $[a, b]$ ,  $f(a) \neq f(b)$ , and  $N$  is between  $f(a)$  and  $f(b)$  then there exists  $c \in (a, b)$  that satisfies  $f(c) = N$ .

## Limits

- $\lim_{x \rightarrow a} f(x)$  exists if and only if  $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x)$
- $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$
- $\lim_{\theta \rightarrow 0} \frac{1 - \cos \theta}{\theta} = 0$

## Derivatives

- $f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$
- $(\cot x)' = -\csc^2 x$
- $(\csc x)' = -\csc x \cdot \cot x$

## Trigonometric

- $\sin^2 \theta + \cos^2 \theta = 1$
- $\sin(2\theta) = 2 \sin \theta \cos \theta$
- $\cos(2\theta) = \cos^2 \theta - \sin^2 \theta$   
 $= 1 - 2 \sin^2 \theta$   
 $= 2 \cos^2 \theta - 1$
- Table of Trig Values

$x$	0	$\pi/6$	$\pi/4$	$\pi/3$	$\pi/2$
$\sin(x)$	0	$1/2$	$\sqrt{2}/2$	$\sqrt{3}/2$	1
$\cos(x)$	1	$\sqrt{3}/2$	$\sqrt{2}/2$	$1/2$	0
$\tan(x)$	0	$\sqrt{3}/3$	1	$\sqrt{3}$	DNE